2.1. Trigonometric Integrals

Some of the following trigonometry identities may be needed.

- A.) $\cos^2 x + \sin^2 x = 1$
- B.) $\sin 2x = 2 \sin x \cos x$
- C.) $\cos 2x = 2\cos^2 x - 1$ so that $\cos^2 x = \dfrac{1 + \cos 2x}{2}$
- D.) $\cos 2x = 1 - 2\sin^2 x$ so that $\sin^2 x = \dfrac{1 - \cos 2x}{2}$
- E.) $\cos 2x = \cos^2 x - \sin^2 x$
- F.) $1 + \tan^2 x = \sec^2 x$ so that $\tan^2 x = \sec^2 x - 1$
- G.) $1 + \cot^2 x = \csc^2 x$ so that $\cot^2 x = \csc^2 x - 1$

2.2 The direct integration for trig. Functions

- 1.) $\int \cos x \, dx = \sin x + C$
- 2.) $\int \sin x \, dx = -\cos x + C$
- 3.) $\int \sec^2 x \, dx = \tan x + C$
- 4.) $\int \csc^2 x \, dx = -\cot x + C$
- 5.) $\int \sec x \tan x \, dx = \sec x + C$
- 6.) $\int \csc x \cot x \, dx = -\csc x + C$
- 7.) $\int \tan x \, dx = \ln|\sec x| + C$
- 8.) $\int \cot x \, dx = \ln|\sin x| + C$
- 9.) $\int \sec x \, dx = \ln|\sec x + \tan x| + C$
- 10.) $\int \csc x \, dx = \ln|\csc x - \cot x| + C$

EXAMPLE Evaluate

$$\int \sin^3 x \cos^2 x \, dx.$$

Solution This is an example of Case 1.

$$\begin{aligned}
\int \sin^3 x \cos^2 x \, dx &= \int \sin^2 x \cos^2 x \sin x \, dx && m \text{ is odd.} \\
&= \int (1 - \cos^2 x) \cos^2 x \, (-d(\cos x)) && \sin x \, dx = -d(\cos x) \\
&= \int (1 - u^2)(u^2)(-du) && u = \cos x \\
&= \int (u^4 - u^2) \, du && \text{Multiply terms.} \\
&= \frac{u^5}{5} - \frac{u^3}{3} + C = \frac{\cos^5 x}{5} - \frac{\cos^3 x}{3} + C. && \blacksquare
\end{aligned}$$

EXAMPLE Evaluate

$$\int \cos^5 x \, dx.$$

Solution This is an example of Case 2, where $m = 0$ is even and $n = 5$ is odd.

$$\begin{aligned}
\int \cos^5 x \, dx = \int \cos^4 x \cos x \, dx &= \int (1 - \sin^2 x)^2 \, d(\sin x) && \cos x \, dx = d(\sin x) \\
&= \int (1 - u^2)^2 \, du && u = \sin x \\
&= \int (1 - 2u^2 + u^4) \, du && \text{Square } 1 - u^2. \\
&= u - \frac{2}{3}u^3 + \frac{1}{5}u^5 + C = \sin x - \frac{2}{3}\sin^3 x + \frac{1}{5}\sin^5 x + C. && \blacksquare
\end{aligned}$$

EXAMPLE Evaluate

$$\int \sin^2 x \cos^4 x \, dx.$$

Solution This is an example of Case 3.

$$\int \sin^2 x \cos^4 x \, dx = \int \left(\frac{1 - \cos 2x}{2}\right)\left(\frac{1 + \cos 2x}{2}\right)^2 dx \qquad m \text{ and } n \text{ both even}$$

$$= \frac{1}{8}\int (1 - \cos 2x)(1 + 2\cos 2x + \cos^2 2x)\, dx$$

$$= \frac{1}{8}\int (1 + \cos 2x - \cos^2 2x - \cos^3 2x)\, dx$$

$$= \frac{1}{8}\left[x + \frac{1}{2}\sin 2x - \int (\cos^2 2x + \cos^3 2x)\, dx\right].$$

For the term involving $\cos^2 2x$, we use

$$\int \cos^2 2x \, dx = \frac{1}{2}\int (1 + \cos 4x)\, dx$$

$$= \frac{1}{2}\left(x + \frac{1}{4}\sin 4x\right). \qquad \text{Omitting the constant of integration until the final result}$$

For the $\cos^3 2x$ term, we have

$$\int \cos^3 2x \, dx = \int (1 - \sin^2 2x)\cos 2x \, dx \qquad u = \sin 2x, \; du = 2\cos 2x \, dx$$

$$= \frac{1}{2}\int (1 - u^2)\, du = \frac{1}{2}\left(\sin 2x - \frac{1}{3}\sin^3 2x\right). \qquad \text{Again omitting } C$$

Combining everything and simplifying, we get

$$\int \sin^2 x \cos^4 x \, dx = \frac{1}{16}\left(x - \frac{1}{4}\sin 4x + \frac{1}{3}\sin^3 2x\right) + C. \qquad \blacksquare$$

EXAMPLE Evaluate

$$\int_0^{\pi/4} \sqrt{1 + \cos 4x}\, dx.$$

Solution To eliminate the square root, we use the identity

$$\cos^2 \theta = \frac{1 + \cos 2\theta}{2} \quad \text{or} \quad 1 + \cos 2\theta = 2\cos^2 \theta.$$

With $\theta = 2x$, this becomes

$$1 + \cos 4x = 2\cos^2 2x.$$

Therefore,

$$\int_0^{\pi/4} \sqrt{1 + \cos 4x}\, dx = \int_0^{\pi/4} \sqrt{2\cos^2 2x}\, dx = \int_0^{\pi/4} \sqrt{2}\sqrt{\cos^2 2x}\, dx$$

$$= \sqrt{2}\int_0^{\pi/4} |\cos 2x|\, dx = \sqrt{2}\int_0^{\pi/4} \cos 2x\, dx \qquad \cos 2x \geq 0 \text{ on } [0, \pi/4]$$

$$= \sqrt{2}\left[\frac{\sin 2x}{2}\right]_0^{\pi/4} = \frac{\sqrt{2}}{2}[1 - 0] = \frac{\sqrt{2}}{2}.$$ ■

2.3 integrals of Power of tan x and sec x

We know how to integrate the tangent and secant and their squares. To integrate higher powers, we use the identities $\tan^2 x = \sec^2 x - 1$ and $\sec^2 x = \tan^2 x + 1$, and integrate by parts when necessary to reduce the higher powers to lower powers.

EXAMPLE : Evaluate

$$\int \tan^4 x\, dx.$$

Solution

$$\int \tan^4 x\, dx = \int \tan^2 x \cdot \tan^2 x\, dx = \int \tan^2 x \cdot (\sec^2 x - 1)\, dx$$

$$= \int \tan^2 x \sec^2 x\, dx - \int \tan^2 x\, dx$$

$$= \int \tan^2 x \sec^2 x\, dx - \int (\sec^2 x - 1)\, dx$$

$$= \int \tan^2 x \sec^2 x\, dx - \int \sec^2 x\, dx + \int dx.$$

$$u = \tan x, \qquad du = \sec^2 x\, dx$$

and have

$$\int u^2\, du = \frac{1}{3}u^3 + C_1.$$

The remaining integrals are standard forms, so

$$\int \tan^4 x\, dx = \frac{1}{3}\tan^3 x - \tan x + x + C. \quad \blacksquare$$

EXAMPLE Evaluate

$$\int \sec^3 x\, dx.$$

Solution We integrate by parts using

$$u = \sec x, \qquad dv = \sec^2 x\, dx, \qquad v = \tan x, \qquad du = \sec x \tan x\, dx.$$

Then

$$\begin{aligned}\int \sec^3 x\, dx &= \sec x \tan x - \int (\tan x)(\sec x \tan x\, dx) \\ &= \sec x \tan x - \int (\sec^2 x - 1)\sec x\, dx \qquad \tan^2 x = \sec^2 x - 1 \\ &= \sec x \tan x + \int \sec x\, dx - \int \sec^3 x\, dx.\end{aligned}$$

Combining the two secant-cubed integrals gives

$$2\int \sec^3 x\, dx = \sec x \tan x + \int \sec x\, dx$$

and

$$\int \sec^3 x\, dx = \frac{1}{2}\sec x \tan x + \frac{1}{2}\ln|\sec x + \tan x| + C. \quad \blacksquare$$

2.4 Products of sines and cosines

The integrals

$$\int \sin mx \sin nx\, dx, \qquad \int \sin mx \cos nx\, dx, \qquad \text{and} \qquad \int \cos mx \cos nx\, dx$$

arise in many applications involving periodic functions. We can evaluate these integrals through integration by parts, but two such integrations are required in each case. It is simpler to use the identities

$$\sin mx \sin nx = \frac{1}{2}[\cos(m-n)x - \cos(m+n)x],$$

$$\sin mx \cos nx = \frac{1}{2}[\sin(m-n)x + \sin(m+n)x],$$

$$\cos mx \cos nx = \frac{1}{2}[\cos(m-n)x + \cos(m+n)x].$$

EXAMPLE Evaluate

$$\int \sin 3x \cos 5x\, dx.$$

Solution From Equation with $m = 3$ and $n = 5$, we get

$$\int \sin 3x \cos 5x\, dx = \frac{1}{2}\int [\sin(-2x) + \sin 8x]\, dx$$

$$= \frac{1}{2}\int (\sin 8x - \sin 2x)\, dx$$

$$= -\frac{\cos 8x}{16} + \frac{\cos 2x}{4} + C.$$

Homework solve ; Ex.8.2 page 466

1,3,5,9,12,18,19,27,31,40,44,51,55,68